

Caustinerf arsenical 6.5g



Composition:

Ephedrine hydrochloride 1.00g, Lidocaine 30.00g, Arsenic trioxide 30.00g

Directions For Use

Posology:

The medical product is applied as a pellet of paste, the size of a pinhead (about 1 mm in diameter), **not more than 10mg**.

Method and route of administration:

Local use only:

Using an excavator, remove the softened dentin and get as near as possible to the pulp.

Pulp removal is slower when the pulp remains protected by a thick layer of dentin and when dentin is hard. Apply a pellet of paste to the cavity floor without exerting excessive pressure and beginning onto the walls. Cover with a temporary cement of relatively soft consistency, so as not to exert too much pressure, which could cause an arsenical "flare-up".

Re-open the tooth no later than 7 days after.



Special warning:

It is imperative to handle the paste very carefully and to observe the usual precautions regarding arsenical products, i.e.:

- Never apply a pellet weighing more than 10 gm of finished product.
- The pulp cavity should be tightly sealed with a temporary occlusive, non compressive cement so as to avoid any risk of arsenical "flare-up".
- Not to leave the pellet in place for more than 7 consecutive days.
- After 7 days, the tooth must be reopened and the entire remaining product must be removed (the product is stained blue by ultramarine blue).
- Clean the pulp cavity after pulpectomy and check before permanent filling that there is no trace of arsenical product left.

Cost Per Application: Approx. Rs.5/-